

High resolution spectroscopy of the high velocity hot post–AGB star LS III +52°24 (IRAS 22023+5249) ★

G. Sarkar¹, M. Parthasarathy², and P. García–Lario³

¹ Department of Physics, Indian Institute of Technology, Kanpur–208016, U.P., India
e-mail: gsarkar@iitk.ac.in

² Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman,
Oklahoma 73019, USA
e-mail: partha@nhn.ou.edu

³ Herschel Science Centre, Research and Scientific Support Department of ESA, European Space
Astronomy Centre (ESAC), Villafranca del Castillo, P.O. Box 50727,E–28080 Madrid, Spain
e-mail: Pedro.Garcia-Lario@sciops.esa.int

Received / Accepted

ABSTRACT

Aims. To investigate the first high resolution optical spectrum of the B–type star, LS III +52°24, identified as the optical counterpart of the hot post–AGB candidate IRAS 22023+5249 (I22023).

Methods. We carried out detailed identifications of the observed absorption and emission features in the high resolution spectrum (4290 – 9015Å) of I22023 obtained with the Utrecht Echelle Spectrograph on the 4.2m William Herschel Telescope. Using Kurucz’s WIDTH9 program and the spectrum synthesis code, SYNSPEC, we determined the atmospheric parameters and abundances. The photospheric abundances were derived under the LTE approximation. The NEBULAR package under IRAF was used to estimate the electron temperature (T_e) and the electron density (N_e) from the [N II] and [S II] lines.

Results. We estimated $T_{\text{eff}}=24000$ K, $\log g=3.0$, $\xi_t=7$ kms^{−1}. The derived CNO abundances suggest an evolved star with $C/O < 1$. P–Cygni profiles of hydrogen and helium indicate ongoing post–AGB mass loss. The presence of [N II] and [S II] lines and the non–detection of [O III] indicate that photoionisation has just started. The derived nebular parameters ($T_e=7000$ K, $N_e=1.2\times10^4$ cm^{−3}) are comparable to those measured in young compact planetary nebulae (PNe). The nebular expansion velocity was estimated to be 17.5 kms^{−1}.

Conclusions. The observed spectral features, large heliocentric radial velocity (-148.31 ± 0.60 kms^{−1}), atmospheric parameters and chemical composition indicate that I22023, at a distance of 1.95 kpc, is an evolved post–AGB star belonging to the old disk population. The nebular parameters suggest that the central star may be evolving into a compact, young PN, similar to Hen3–1357.

Key words. Stars: AGB and post–AGB – Stars: early–type – Stars: abundances – Stars: evolution

1. Introduction

The discovery of high galactic latitude cool (F, G, K) and hot (OB) post–AGB supergiants, e.g. HD 161796 (Parthasarathy & Pottasch 1986) and LSII +34°26 (Parthasarathy 1993) indicated that K, G, F, A, OB post–AGB supergiants form an evolutionary sequence in the transition region from the tip of the AGB into early stages of planetary nebulae (PNe). Since then, several cool and hot post–AGB candidates have been identified (Pottasch & Parthasarathy 1988, Parthasarathy & Pottasch 1989, García–Lario et al. 1997a; Parthasarathy et al. 2000a). Gauba & Parthasarathy (2003, 2004) analysed the UV spectra and circumstellar dust envelopes of several hot post–AGB stars from the above lists, including LS III +52°24. High resolution optical spectra of only a few hot post–AGB stars have been studied (Sarkar et al. 2005 and references therein). To ascertain their evolutionary nature, atmospheric parameters and chemical compositions, there is a clear need to study a bigger sample of these objects.

The optically bright B–type star, LS III +52°24 identified with the IR source IRAS 22023+5249 (hereafter, I22023) has far–IR colors similar to PNe (Table 1). It is listed in Wackerling’s (1970) Catalog of early–type emission–line stars. In this paper, we present the first high resolution optical spectrum of I22023. Recent, ground–based, high resolution, near–IR images, have shown H₂ emission arising close to the central star, possibly in a bipolar morphology (Volk et al. 2004). Kelly & Hrivnak (2005) also detected Hydrogen Br _{γ} 2.166 μ m line in emission. Gauba & Parthasarathy (2004) indicated the presence of weak amorphous (10.8 μ m) and crystalline (33.6 μ m) silicate features in the ISO spectrum of the object. However, another look at the ISO spectrum suggests that this identification was possibly in error. The ISO spectrum of I22023 is very noisy and displays only a featureless continuum with no 10.8 μ m and 33.6 μ m features.

Table 1. Details of the star

IRAS	Name	RA 2000	DEC 2000	l	b	Sp. Type Optical	V	B–V	IRAS Fluxes (Jy)			
									12 μ m	25 μ m	60 μ m	100 μ m
22023+5249	LSIII +52 24	22:04:12.30	+53:04:01.4	99.30	–1.96	B ^a	12.52 ^b	0.69 ^b	1.02	24.69	14.52	3.93L

^aSpectral type is from the SIMBAD database. ^bPhotometry is from Hog et al. (2000). L flag indicates that the quoted IRAS flux density is an upper limit.

2. Observations

I22023 was observed on 14 July 2001 using the Utrecht Echelle Spectrograph (UES) on the 4.2m William Herschel Telescope (WHT) at the Roque de los Muchachos Observatory on La Palma. Observations were made with the 31.6 lines/mm echelle grating (E31), SITe1 CCD (2048 $\times$ 2048 pixels of 24 μ m), a slit width of 1'' on the sky and central wavelength of 5500Å resulting in a resolving power of $R \sim 52000$. The wavelength coverage was 4290–4735Å, 4760–5553Å and 5607–9015Å. A Th–Ar comparison lamp was used for wavelength calibration.

Send offprint requests to: G. Sarkar

* Based on observations obtained with the 4.2m William Herschel Telescope and the Utrecht Echelle Spectrograph of the Roque de Los Muchachos Observatory, La Palma, Spain

3. Analysis

The spectrum was extracted using standard reduction procedures in the IRAF package. The reduction steps included overscan correction, trimming, bias subtraction, flat field correction, correction for scattered light and wavelength calibration. The reduced spectrum was continuum-normalised. The final S/N varied from 30 in the blue to 60 towards the red end of the spectrum.

Equivalent widths (W_λ) of the absorption and emission lines were measured. Deblending was done whenever required to obtain Gaussian fits to the blended line profiles. The complete continuum-normalised spectrum of I22023 (Fig. 1) is presented in the appendix. This spectrum would be useful for future observers since post-AGB stars show both short and long-term variability in the absorption and emission line strengths and profiles. P-Cygni profiles detected in these stars are also expected to vary as the stellar wind and post-AGB mass loss rates may show variation as the star evolves. The line identifications (Tables 2 – 5) are based on the Moore multiplet table (1945) and the linelists of Parthasarathy et al. (2000b), Klochkova et al. (2002) and Sarkar et al. (2005). Unidentified lines are denoted by “UN”. Night sky emission lines denoted by “atmos.” were identified from Osterbrock et al. (1996). The laboratory wavelengths, $\log(gf)$ values and excitation potentials (χ) have been extracted from the Kurucz (CD-ROM 23) linelist GFALL. Ivan Hubeny and Thierry Lanz have compiled the Kurucz linelists with improved oscillator strengths from the NIST Atomic Spectra Database. For wavelengths below 7500 Å we have used their data which may be retrieved from <http://nova.astro.umd.edu/Synspec43/synspec-frames-data.html>.

3.1. Description of the spectrum

The absorption and emission lines in the spectrum of I22023 are similar to those detected in other hot post-AGB stars (Parthasarathy et al. 2000b, Klochkova et al. 2002 and Sarkar et al. 2005). In addition to the O I triplet, absorption lines of He I, C II, C III, N II, O II, Ne I, Mg II, Al III, Si III were identified. Emission lines of C II, N II, O I, [O I], Al III, Si II, Fe I, Fe II, [Fe II] and Fe III were also identified in the spectrum of the star. The NaI D1 and D2 lines show a complex structure. The presence of low excitation nebular lines of [N II] and [S II] and the absence of [O III] 5007Å suggest that photoionization has just started, shock excitation may also be playing a role. Balmer lines of H_α , H_β and H_γ show P-Cygni profiles indicating ongoing post-AGB mass loss. Some He I, C II and Fe III lines were also found to have P-Cygni profiles.

3.2. Reddening

For the derived effective temperature (T_{eff}) and gravity of the star (Sec. 3.8), we may assume a B1III spectral type with intrinsic B–V $((B-V)_0) = -0.26$ (Schmidt-Kaler 1982). The observed B–V for I22023 is 0.69 (Hog et al. 2000). The total $E(B-V)$ towards the star is therefore estimated to be 0.95. At the low galactic latitude ($b = -1.96$) of the star, the Diffuse Infrared Background Experiment (DIRBE)/IRAS dust maps (Schlegel et al. 1998) do not give reliable estimates of the interstellar extinction. The interstellar $E(B-V)$ was estimated to be 0.63 based on the strength of the DIB at 5780.410Å (Sec. 3.6). This suggests that I22023 suffers from significant circumstellar reddening. The UV spectrum of the star obtained with the International Ultraviolet Explorer (IUE)

Table 2. Absorption lines in IRAS 22023+5249

$\lambda_{\text{obs.}}$ (Å)	$\lambda_{\text{lab.}}$ (Å)	Ident.	W_λ (Å)	log (gf)	χ (eV)	$ \Delta\lambda $ (Å)	V_r km s ⁻¹
4314.874	4317.139	OII(2)	0.1609	-0.386	22.95–25.82	2.265	-142.53
4317.277	4319.630	OII(2)	0.2197	-0.380	22.96–25.83	2.353	-148.55
4343.245	4345.560	OII(2)	0.1448	-0.346	22.96–25.81	2.315	-144.95
4345.141	4347.413	OII(16)	0.0480	0.024	25.64–28.49	2.272	-141.91
4347.137	4349.426	OII(2)	0.2489	0.060	22.98–25.83	2.289	-143.01
4348.903	4351.260	OII(16)	0.0483	0.227	25.64–28.49	2.357	-147.64
4364.653	4366.895	OII(2)	0.1130	-0.348	22.98–25.82	2.242	-139.15
4385.301	4387.929	HeI(51)	0.3722	-0.883	21.20–24.03	2.628	-164.80
4412.525	4414.899 ^a	OII(5)		0.172	23.42–26.23		blend
4414.702	4416.975 ^b	OII(5)	0.1319	-0.077	23.40–26.21	2.384	blend
4435.158	4437.551	HeI(50)	0.0786	-2.034	21.20–23.99	2.393	-146.91
4478.638	4479.885	AlIII(8)	0.0524	0.900	20.77–23.53		
	+ 4479.971	AlIII(8)		1.020	20.77–23.53		
	+ 4481.126	MgII(4)		0.740	8.86–11.62	2.488	-151.70
4550.109	4552.622	SiIII(2)	0.3600	0.181	19.00–21.72	2.513	-150.73
4565.371	4567.840	SiIII(2)	0.3225	-0.039	19.00–21.72	2.469	-147.28
4572.278	4574.757	SiIII(2)	0.1638	-0.509	19.00–21.71	2.479	-147.70
4588.513	4590.974	OII(15)	0.1206	0.350	25.64–28.34	2.461	-145.95
4593.736	4596.177	OII(15)	0.1507	0.200	25.64–28.34	2.441	-144.46
4627.829	4630.539	NII(5)	0.0983	0.094	18.47–21.14	2.710	-160.70
4636.365	4638.856	OII(1)	0.1682	-0.332	22.95–25.62	2.491	-146.23
4639.279	4641.810	OII(1)	0.3179	0.055	22.96–25.63	2.531	-148.71
4644.997	4647.418	CIII(1)	0.1181	0.070	29.51–32.18	2.421	-141.41
4646.569	4649.135	OII(1)	0.3387	0.308	22.98–25.65	2.566	-150.71
4648.297	4650.16	CIII(1)	0.2709				blend
	+ 4650.841	OII(1)					
	+ 4651.35	CIII(1)					
4659.126	4661.632	OII(1)	0.2431	-0.278	22.96–25.62	2.506	-146.40
4671.265	4673.733	OII(1)	0.0644	-1.090	22.96–25.61	2.468	-143.55
4673.706	4676.235	OII(1)	0.1598	-0.394	22.98–25.63	2.529	-147.38
4693.804	4696.353	OII(1)	0.0346	-1.380	22.98–25.62	2.558	-148.53
4696.824	4699.218	OII(25)	0.0944	0.270	26.21–28.84	2.394	-137.96
4702.873	4705.346	OII(25)	0.0621	0.477	26.23–28.86	2.473	-142.80
4817.027	4819.712	SiIII(9)	0.0556	0.750	25.96–28.53	2.685	-152.26
4826.346	4828.951	SiIII(9)	0.0344	1.090	25.97–28.53	2.605	-146.97
4904.197	4906.830 ^c	OII(28)		-0.161	26.29–28.81	2.633	-146.11
4918.616	4920.35	HeI(49)	0.4414		21.13–23.64		blend
	+ 4921.931	HeI(48)		-0.435	21.20–23.72		
4921.878	4924.529	OII(28)	0.0851	0.074	26.29–28.80	2.651	-146.63

^aOII(5) 4414.899Å is blended with FeII(32) Å emission. ^bOII(5) 4416.975Å is blended with the absorption component of FeIII(114) 449.596 P–Cygni profile. ^cOII(28) 4906.88 Å is blended with [FeII](20F) 4905.35 Å emission.

Table 2. Absorption lines in IRAS 22023+5249

$\lambda_{\text{obs.}}$ (Å)	$\lambda_{\text{lab.}}$ (Å)	Ident.	W_{λ} (Å)	$\log(gf)$	χ (eV)	$ \Delta\lambda $ (Å)	V_r km s ⁻¹
4940.268	4943.005	OII(33)	0.0825	0.239	26.54–29.05	2.737	–151.24
5000.141	5002.703	NII(4)	0.0328	–1.022	18.45–20.92	2.562	–138.77
5002.414	5005.150	NII(19)	0.0336	0.594	20.65–23.13	2.736	–149.12
5007.895	5010.621	NII(4)	0.0851	–0.607	18.45–20.92	2.726	–148.34
5042.253	5044.8	CII(35)	0.1148				blend
	+ 5045.098	NII(4)					
5044.65	5047.2	CII(35)	0.0825				blend
	+ 5047.736	HeI(47)					
5130.277	5132.947	CII(16)	0.0376	–0.211	20.69–23.10		blend
	+ 5133.282	CII(16)		–0.178	20.69–23.10		
5140.722	5143.495	CII(16)	0.0640	–0.212	20.69–23.10		blend
5142.407	5145.165	CII(16)	0.0523	0.189	20.70–23.10		blend
5148.32	5151.085	CII(16)	0.0446	–0.179	20.70–23.10	2.765	–146.16
5216.486		UN	0.0471				
5486.97	5487.43	DIB	0.0343			0.460	–10.28
5663.356	5666.630	NII(3)	0.1400	–0.045	18.45–20.64	3.274	–158.46
5672.88	5676.02	NII(3)	0.1433	–0.367	18.45–20.63	3.14	–151.09
5676.262	5679.56	NII(3)	0.2223	0.250	18.47–20.65	3.298	–159.33
5682.922	5686.21	NII(3)	0.0382	–0.549	18.45–20.63	3.288	–158.60
5693.475	5695.920	CIII(2)	0.1418	0.017	32.08–34.26		
	+ 5696.604	AlIII(2)		0.230	15.63–17.81	3.129	–149.91
5707.504	5710.770	NII(3)	0.0552	–0.518	18.47–20.64	3.266	–156.70
5719.637	5722.730	AlIII(2)	0.0786	–0.070	15.63–17.80	3.093	–147.27
5736.688	5739.734	SiIII(4)	0.2575	–0.160	19.71–21.87	3.046	–144.34
5779.728	5780.410	DIB	0.3405			0.682	–20.52
5796.444	5797.030	DIB	0.0440			0.586	–15.46
6139.673	6143.063	NeI(1)	0.0282	–0.350	16.61–18.62	3.39	–150.68
6195.200	6195.990	DIB	0.0756			0.790	–23.38
6202.191	6203.060	DIB	0.0966			0.869	–27.16
6269.073		UN	0.0691				
6282.734	6283.86 ^d	DIB	0.7319			1.126	–38.89
6398.61	6402.246	NeI(1)	0.0543	0.360	16.61–18.54	3.636	–155.48
6612.857	6613.630	DIB	0.1268			0.773	–20.19
6637.522	6641.031	OII(4)	0.0238	–0.884	23.40–25.27	3.509	–143.64
6717.943	6721.388	OII(4)	0.0963	–0.610	23.43–25.27	3.445	–138.89
7766.623	7771.944	OI(1)	0.576	0.320	9.14–10.74		blend
7769.416	7774.166	OI(1)	0.434	0.170	9.14–10.74		blend
	+ 7775.388	OI(1)		–0.050	9.14–10.74		
8592.524		UN	0.265				
8646.868	8648.28	DIB	0.466			1.412	–34.11

^dDIB 6283.86Å is blended with telluric absorption lines in this region.

Table 3. Emission lines in IRAS 22023+5249

$\lambda_{\text{obs.}}$ (Å)	$\lambda_{\text{lab.}}$ (Å)	Ident.	W_{λ} (Å)	$\log(gf)$	χ (eV)	$ \Delta\lambda $ (Å)	V_r km s ⁻¹
4411.134	4413.601 ^a	FeII(32)	0.0600	-3.870	2.67–5.48		blend
4811.856	4814.55	[FeII](20F)	0.0409			2.694	-153.00
4902.59	4905.35 ^b	[FeII](20F)	0.0287				blend
5038.313	5041.024	SiII(5)	0.1095	0.291	10.06–12.52	2.711	-146.47
5053.306	5055.984	SiII(5)	0.2758	0.593	10.07–12.52		blend
	+ 5056.317	SiII(5)		-0.359	10.07–12.52		
5088.76		UN	0.0244				
5155.834	5158.81	[FeII](19F)	0.0985				blend
5191.418	5193.909	FeIII(5)	0.0594	-2.852	8.65–11.04		blend
	+ 5194.384	FeIII(5)			8.65–11.04		
5195.026	5197.929	FeI(1091)	0.1135	-1.640	4.30–6.68	2.903	-152.68
5197.427		UN	0.084				
5232.982		UN	0.0412				
5240.659	5243.306	FeIII(113)	0.0997	0.405	18.26–20.62		blend
	+ 5243.773	FeI(1089)		-1.150	4.25–6.62		
5258.712	5261.61	[FeII](19F)	0.0520			2.898	-150.36
5270.327	5273.38	[FeII](18F)	0.0481			3.053	-158.81
5273.648		UN	0.0487				
5279.665		UN	0.0756				
5282.071		UN	0.0277				
5286.85		UN	0.0632				
5296.105	5299.044	OI(26)	0.1163	-2.140	10.98–13.32		blend
5297.325		UN	0.0366				
5299.853		UN	0.0368				
5751.43	5754.8	[NII](3F)	0.0436				weak
5830.998	5834.06	FeII(165)	0.0705	-3.738	5.57–7.69	3.062	-142.58
5917.114	5920.124	FeIII(115)	0.1332	-0.034	18.78–20.87	3.01	-137.66
5926.609	5929.685	FeIII(114)	0.0964	0.351	18.50–20.59	3.076	-140.75
5950.424	5953.613	FeIII(115)	0.0989	0.186	18.78–20.86	3.226	-147.69
5954.336	5957.559	SiII(4)	0.1449	-0.301	10.06–12.14	3.223	-147.43
5975.851	5978.930	SiII(4)	0.4134	0.004	10.07–12.14	3.079	-139.62
5996.369	5999.70	AlII(93)	0.1843		15.52–17.57		blend
	+ 5999.83	AlII(93)			15.52–17.57		
6029.382	6032.67	FeI(1082)	0.2320		4.20–6.25	3.288	-148.64

^aFeII(32) 4413.601Å is blended with OII(5) 4414.899Å absorption feature. ^b[FeII](20F) 4905.35Å is blended with OII(28) 4906.88Å absorption feature.

satellite also indicates considerable circumstellar reddening $(E(B-V))_{\text{circumstellar}} = 0.44$, Gauba & Parthasarathy 2003).

3.3. Distance estimates

Assuming I22023 to be a normal population I star, we may adopt the simple approximation given in Lang (1980) for the velocity structure of the Galactic disk which expresses the heliocentric radial velocity as a function of distance and Galactic coordinates of the star :

$$v_{\odot} = -\delta v_{\odot} + A D \sin 2l \cos^2 b \text{ [kms}^{-1}\text{]},$$

where, $v_{\odot}$ is the heliocentric radial velocity of I22023; $\delta v_{\odot}$ is the correction term to convert from heliocentric radial velocity to velocity in the Local Standard of Rest (v_{lsr}) and is 13.19 kms^{-1} for I22023; Oort's constant, $A \simeq 15 \text{ km s}^{-1} \text{ kpc}^{-1}$; D is the distance to the star in kpc and the Galactic coordinates for I22023, $(l; b) = (99.30; -1.96)$. Using $v_{\odot} = -148.31 \text{ km}^{-1}$ (Sec. 3.4), we derive a distance of 28.2 kpc to I22023. Such a large distance is unphysical and suggests that I22023 lies beyond the edge of our Galaxy. It is therefore, unlikely that I22023 is a normal population I B giant/supergiant star.

On the post-AGB evolutionary tracks of Schönberner (1987), a post-AGB star with $T_{\text{eff}} = 24000\text{K}$ and $\log g = 3.0$ (Sec. 3.8) has a core mass, $M_c = 0.565 M_{\odot}$. If I22023 is a post-AGB star, then using the relation between core-mass and quiescent luminosity maximum (Wood & Zarro 1981) for AGB stars, we estimate $M_{\text{bol}} = -4.30$. Applying the bolometric correction ($= -2.43$) for a B1III star (Lang 1992) we derive $M_v = -1.87$. The observed V magnitude ($m_v = 12.52$; Hog et al. 2000) of the star was corrected for extinction using $E(B-V) = 0.95$ ($A_v = 2.94$). Distance modulus method then yields a distance of 1.95 kpc to the star.

3.4. Radial velocity

Heliocentric radial velocities have been derived from wavelength shifts of the well defined absorption and emission lines (Tables 2, 3, 4 and 5). The mean heliocentric radial velocities from the absorption and emission lines (Tables 2 and 3) are $-148.31 \pm 0.60 \text{ kms}^{-1}$ and $-144.13 \pm 0.72 \text{ kms}^{-1}$ respectively. Radial velocity measurements of the forbidden lines have been excluded in estimating the mean. The quoted errors refer to the probable errors of estimation. Fig.2 shows the radial velocity trends with respect to the equivalent widths (W_{λ}) and lower excitation potentials (LEP) of the absorption and emission lines respectively.

The mean heliocentric radial velocity of the [N II], [O I] and [Fe II] lines is $-152.90 \pm 0.96 \text{ kms}^{-1}$. [S II] 6717.0Å and 6731.3Å lines in I22023 have a markedly different heliocentric velocity corresponding to $-171.93 \pm 1.36 \text{ kms}^{-1}$. The different radial velocities argue for a non-spherical nebula.

3.5. Wind velocities and mass loss rate from the P-Cygni profiles

The Balmer lines, H_{α} , H_{β} , H_{γ} , a few He I, CII and Fe III lines in I22023 show P-Cygni behaviour (Fig. 1, Table 4) indicating ongoing mass-loss. Wind velocities were estimated from the blue absorption edges of the well defined and unblended P-Cygni profiles (Table 4). The wind velocities of the Fe III(5) lines are markedly different from those of the other species. However, the wind velocities do not show any obvious trend with the lower excitation potentials or ionization potentials of the species. The mean wind velocity from the Balmer, He I and C II lines is 187.48 kms^{-1} and that from the Fe III(5) lines is 149.26 kms^{-1} . This difference in velocities indicates a deviation




Fig. 2. Radial velocity trends of the absorption and emission lines in IRAS22023+5249. Radial velocity measurements of the forbidden lines have not been plotted.

from spherical symmetry – possibly a bipolar morphology and the presence of a dense equatorial torus (e.g. Welch et al. 1999; Sahai et al. 2005). HST images may reveal the true morphology of the object.

The Balmer lines, H_α , H_β and H_γ in I22023 are shown in Fig. 3. The velocities are in the heliocentric frame. Detailed modelling of these lines to derive the mass loss rate would be the subject of a future paper. The equivalent widths of the H_α emission components are related to the mass loss rates in OB stars (Leitherer 1988). The H_α emission component in I22023 has an equivalent width ($W_\lambda = 37.8\text{\AA}$) comparable to that of the O8I star, HD152408 ($W_\lambda = 34.7\text{\AA}$). From this, we estimate a mass loss rate of $1.23 \times 10^{-5} M_\odot \text{yr}^{-1}$ (Leitherer 1988). However, this value may be treated with caution. In I22023, the large equivalent width of the H_α emission component may be due to a large amount of gas in its circumstellar envelope or the presence of a bipolar envelope (Volk et al. 2004) and may not be directly related to the mass loss rate.

3.6. Diffuse interstellar bands (DIBs)

DIBs $5487.43\text{\AA}$, $5780.410\text{\AA}$, $5797.030\text{\AA}$, $6195.990\text{\AA}$, $6203.060\text{\AA}$, $6283.86\text{\AA}$, $6613.630\text{\AA}$ and $8648.28\text{\AA}$ were identified in the spectrum of the star (Table 2). They have also been identified in




Fig. 3. Observed Balmer profiles (dotted) in I22023 show P Cygni behaviour. The velocities are in the heliocentric frame.

other hot post-AGB stars e.g. IRAS01005+7910 (Klochova et al. 2002), IRAS13266-5551 & IRAS17311-4924 (Sarkar et al. 2005).

DIBs may have their origin in both the interstellar and the circumstellar media. A lower and upper limit on the predicted velocities of interstellar lines may be obtained by substituting $D = 0$ and $D = 1.95$ kpc (the assumed distance to I22023) in the formula stated in Sec. 3.3 above (see e.g. Bakker et al. 1996). This yields a range of -13.19 km s^{-1} to -22.53 km s^{-1} . This may however be treated as only a rough estimate.

A comparison of the heliocentric radial velocities of the identified DIBs with the estimate above suggests that all of these lines most likely originate in the interstellar medium. From the strength of the band at $5780.410 \text{ \AA}$ ($W_\lambda = 340.5 \text{ m\AA}$), we estimate interstellar $E(B-V) \approx 0.63$ (Herbig 1993).

3.7. Na I D_2 and Na I D_1 lines

Six components were identified in the Na I D_2 and Na I D_1 lines (Fig. 4, Table 5). The velocities of absorption component 1 and emission component 2 are comparable with the mean heliocentric radial velocities of the absorption and emission lines in the star (Sec. 3.4) suggesting that component 1 is of photospheric origin and component 2 arises in an extended envelope around the central star. Comparing the heliocentric velocities of absorption components 3, 4 and 5 with those of DIBs observed in the spectrum of the star we may infer that these components originate in the interstellar medium. Component 6 is observed in emission with a velocity very different from the envelope velocity. This component may arise in a disk or in outflows around the central star. The velocity of




Fig. 4. Na I D₂ and Na I D₁ lines in I22023. The various absorption and emission components (Sec. 3.7, Table 5) have been labelled.

this component is comparable with the expansion velocity estimated from the nebular lines (Sec. 3.9.2).

3.8. Atmospheric parameters and abundances from absorption line spectrum

The detection of He I and Si III absorption lines in addition to the C III lines and the absence of He II absorption lines, indicates a B0 – B1 supergiant spectral type for the central star. Comparison of the UV(IUE) spectrum of I22023 with standard stars suggests that the star is a B2-supergiant (Gauba & Parthasarathy 2003). Similar to our analysis of the high resolution optical spectrum of the hot post-AGB star, IRAS13266–5551 (Sarkar et al. 2005), we used Kurucz’s WIDTH 9 program and the spectrum synthesis code, SYNSPEC (Hubeny et al. 1985) along with solar metallicity Kurucz (1994) model atmospheres to derive the atmospheric parameters and elemental abundances of I22023 under the LTE approximation.

The largest number of absorption lines in I22023 are those of O II and N II. We derived the oxygen and nitrogen abundance with the WIDTH 9 program for various combinations of T_{eff} , $\log g$ and ξ_t . We covered $18000 \text{ K} \leq T_{\text{eff}} \leq 25000 \text{ K}$. From the Kurucz (1994) model atmospheres, the $\log g$ value was limited to a minimum of 3.0. For each combination of these parameters, we then synthesised the spectrum using SYNSPEC. The best fit to the observed spectrum was obtained for $T_{\text{eff}} = 24000 \text{ K}$, $\log g = 3.0$, $\xi_t = 7 \text{ km s}^{-1}$. Since strong lines are usually affected by microturbulence, the use of these lines in determining the atmospheric parameters of the star may contribute

to systematic errors. In estimating the atmospheric parameters and abundances, we excluded lines with $W_\lambda \geq 200 \text{ mÅ}$. Line blends were also excluded from this analysis.

3.8.1. He I lines

Several absorption and P-Cygni type He I lines were identified in the spectrum of I22023. The absorption lines are blends with the exception of the strong 4387.929 Å He I(51) ($W_\lambda = 372.2 \text{ mÅ}$) line and the He I(50) 4437.551 Å line ($W_\lambda = 78.6 \text{ mÅ}$). For the derived atmospheric parameters of the star, using spectrum synthesis, we estimated the helium abundance from the He I(50) line.

3.8.2. C II and C III lines

Both C II and C III absorption lines were detected in the spectrum of the star. However, the number of these lines in I22023 is few. Further, all the identified carbon lines are blends with the exception of the CII(16) line at 5151.085 Å . The carbon abundance was therefore estimated using spectrum synthesis alone. The region of the CII(16) lines ($\sim 5132 \text{ Å} - 5151 \text{ Å}$) was used for this purpose.

3.8.3. OII lines and O I triplet

The largest number of absorption lines in the spectrum are those of O II. The O II lines with $W_\lambda \geq 200 \text{ mÅ}$ may be sensitive to NLTE effects. Such strong lines were neglected while estimating the oxygen abundance using WIDTH9. For the derived atmospheric parameters and oxygen abundance, using the SYNSPEC code, we could not obtain a good fit to such lines.

The (total) equivalent width of the O I triplet in the spectrum of I22023 is 1.01 Å . This is comparable to the 0.95 Å equivalent width of the O I triplet in the spectrum of the B1.5Ia hot post-AGB star, LSII+34°26 (García-Lario et al. 1997b; Arkhipova et al. 2001). The O I triplet at 7773 Å is known to be sensitive to NLTE effects. Using $\log \epsilon(\text{O}) = 8.90$ derived from the O II lines in the spectrum of I22023 (Table 6a), we could not obtain a fit to the O I triplet.

3.8.4. N II and Ne I lines

Several N II and two Ne I lines were identified in I22023. The abundances of these lines were estimated using WIDTH9. Once again, the mismatch between the strong ($W_\lambda \geq 200 \text{ mÅ}$) observed and synthetic N II lines indicates the need for a NLTE analysis of the spectrum of this star.

3.8.5. Mg II lines

Only one Mg II line could be identified in the spectrum of the star. This line is blended with Al III(8). Since the Al III abundance in I22023 is uncertain (see below), we did not attempt to estimate the magnesium abundance from the blended 4481.126 Å Mg II(4) line.

3.8.6. Al III lines

Only four Al III lines could be identified. Three of these lines are blends with other species. We therefore, estimated the aluminium abundance from the single 5722.730 Å Al III(2) line using

spectrum synthesis analysis and derived $\log \epsilon(\text{Al})=6.79$. This abundance from a single line with $W_\lambda=78.6\text{m}\text{\AA}$ may be treated as an upper limit.

3.8.7. Si III lines

The derived silicon abundance, $[\text{Si}/\text{H}]=-0.12$ suggests that I22023 may be slightly metal deficient.

3.8.8. Iron lines

Iron abundance could not be estimated since the iron lines in I22023 appear only in emission or show P-Cygni profiles.

3.8.9. Error estimates

The standard deviation (σ) which measures the scatter in the abundances due to individual lines of a species was estimated using WIDTH9 (Table 6a). The true error, $\sigma/\sqrt{n}$, would be smaller for species with a greater number of lines (n).

Table 6b gives the uncertainties in the abundances due to uncertainties in the model atmospheric parameters, taking $\Delta T_{\text{eff}}=-1000\text{K}$, $\Delta \log g=+0.5$ and $\Delta \xi_t=+1 \text{ km s}^{-1}$. The quadratic sum of the uncertainties due to the model parameters is given by σ_m .

3.9. Analysis of the emission line spectrum

Several permitted and forbidden emission lines were identified in the spectrum of the star (Table 3). Nebular parameters and expansion velocities were determined using the forbidden lines.

3.9.1. Nebular parameters

In the absence of a flux calibrated spectrum for I22023, it is not possible to obtain the absolute fluxes in the observed emission lines. However, reliable emission line flux ratios may be deduced by combining the observed equivalent widths (Table 3) with estimates of the stellar continuum flux distribution in the regions of the emission lines. The latter were obtained for the derived atmospheric parameters of the star (Sec. 3.8) using the SYNSEPC code and Kurucz model atmospheres. The emission line fluxes thus estimated are free from the effects of both interstellar and circumstellar reddening.

The $[\text{S II}] \lambda 6717/\lambda 6731$ line ratio is an electron density diagnostic and the $[\text{N II}] (\lambda 6548+\lambda 6583)/\lambda 5755$ line ratio is sensitive to electron temperature. In I22023, the $[\text{N II}] 6583.6\text{\AA}$ emission is blended with the emission component of the C II(2) $6582.88\text{\AA}$ P-Cygni profile. However, comparing the $6582.88\text{\AA}$ C II(2) profile with the $6578.05\text{\AA}$ C II(2) P-Cygni profile (Fig. 1), we may conclude that the contribution of C II(2) to the $[\text{N II}]$ emission profile is negligible. Using the NEBULAR analysis package under IRAF, we obtained T_e Vs. N_e contours for the observed $[\text{S II}]$ and $[\text{N II}]$ diagnostic ratios of 0.5 and 166.7 respectively. From the intersection of the contours we obtained $T_e=7000 \text{ K}$ and $N_e=1.2\times 10^4 \text{ cm}^{-3}$. The high electron density is comparable to that measured in the young, compact PN, Hen3-1357 (Parthasarathy et al. 1993,

Bobrowsky et al. 1998) which evolved from the hot post-AGB stage into a PN in 20–30 yrs. (Parthasarathy et al. 1995).

We could not derive the nebular C, N, O abundances which could then have been compared with the photospheric abundances to estimate the amount of material lost by the star during nebular formation and the chemical composition of the nebula. Such a calculation requires an estimate of the H_β emission line flux. However, H_β in I22023 shows a P-Cygni profile and it is not possible to estimate the nebular emission from this profile.

3.9.2. Expansion velocities

Expansion velocities were estimated from the FWHM of the unblended [OI], [NII] and [SII] lines using $V_{\text{exp}} = 0.50 \text{ FWHM}$ (Table 7). This approximation is valid when emission is confined to a thin spherically symmetric shell. However, I22023 appears to have a bipolar morphology in recent ground based high resolution images (Volk et al. 2004). Further, the observed [OI] 6300.23Å and 6363.88Å line profiles appear to be asymmetric. Though no obvious line split is observed in the weak [OI] lines, their asymmetric nature may indicate the presence of a red and a blue component. This may explain the discrepancy between the expansion velocities estimated from [OI] and [NII] and [SII] lines in I22023, the former being nearly twice that of the latter two species. The mean nebular velocity based on the [NII] and [SII] lines is 17.5 kms^{-1} .

4. Discussion and conclusions

In this paper, we present the first high resolution optical spectrum and detailed line identification of the B-type star, LS III +52°24, identified as the optical counterpart of the hot post-AGB candidate I22023. That I22023 is not a normal population I B star is also suggested by the large heliocentric radial velocity of the star ($-148.31 \pm 0.60 \text{ kms}^{-1}$) measured from the absorption lines in its high resolution optical spectrum. It is more likely that I22023 is a post-AGB star belonging to the old disk population. The presence of absorption lines of He I, C III, Si III alongwith [N II], [O I] and [S II] emission lines indicate a low excitation nebula surrounding an early B-type star. P-Cygni lines of hydrogen, He I, C II and Fe III provide evidence for ongoing post-AGB mass loss.

As a first approximation, using LTE analysis, we estimated $T_{\text{eff}} = 24000 \text{ K} \pm 1000 \text{ K}$, $\log g = 3.0 \pm 0.5$ and $\xi_t = 7 \pm 1 \text{ kms}^{-1}$. The O I triplet and strong He I, O II and some N II absorption lines indicate that NLTE effects are significant in the star. The derived CNO abundances were compared with average CNO abundances for main-sequence B-stars from the Ori OB1 association (Table 6a; Kilian 1992). The CNO abundances suggest that I22023 is an evolved star. We estimated C/O ~ 0.48 . A spectrum with better S/N ratio and NLTE analysis of the absorption lines may provide improved estimates of the atmospheric parameters and abundances. Besides high resolution UV spectra of the object may reveal a greater number of absorption lines in the UV which would help in a more accurate determination of these parameters.

Even though the IRAS cool and hot post-AGB stars show supergiant like spectra indicating an evolutionary sequence in the transition region from the AGB to the PN stage, they seem to show significant differences in chemical composition. In the high galactic latitude hot post-AGB stars severe carbon deficiency is detected indicating that they left the AGB before the third dredge-up (Conlon et al. 1991; McCausland et al. 1992; Moehler & Heber 1998). Similar carbon deficiency

is also detected in the hot post-AGB stars in globular clusters (Moehler et al. 1998; Mooney et al. 2004; Jasiewicz et al. 2004; Thompson et al. 2006). The only exception to this pattern is the field hot post-AGB star, IRAS 01005+7910 which showed an overabundance of carbon (Klochkova et al. 2002). In contrast, amongst IRAS selected cool post-AGB stars, not a single post-AGB star is detected with severe carbon deficiency. The F-type post-AGB stars with 21-micron emission show an overabundance of carbon and s-process elements indicating that they have experienced third dredge-up and evolved from the carbon star stage on the AGB.

Evidence for bipolar nebulosity was seen in ground based H_2 images of I22023 (Volk et al. 2004). Bipolar proto-planetary nebulae (PPNe) are usually attributed to binary systems. In our high resolution optical spectrum, we did not find any evidence for binarity of the hot central star. HST WFPC2 images may provide better insight into the morphology and evolutionary nature of this object.

Far-IR (IRAS) colours similar to PPNe, H_2 emission arising close to the central star with possible bipolar morphology (Volk et al. 2004), large heliocentric radial velocity of $-148.31 \text{ km s}^{-1}$ and absence of large amplitude irregular light variations similar to that observed in luminous blue variables (LBVs) indicate that I22023 cannot be classified as a LBV candidate. I22023 may show low amplitude irregular light variations similar to that observed in SAO 85766, LSII +34°26 and other hot post-AGB stars (Arhipova et al. 2006).

Acknowledgements. GS would like to acknowledge financial support from the Department of Science and Technology (DST), Govt. of India through a grant numbered SR/FTP/PS-67/2005. MP is very thankful to Prof. David Branch, Prof. E. Baron and Prof. Ryan Doezema for their kind support, encouragement and hospitality.

References

- Arkhipova, V.P., Klochkova, V.G., Sokol, G.V., 2001, *AstL*, 27, 99
- Arkhipova, V.P., Ikonnikova, N.P., Komissarova, G.V., Noskova, R.I. 2006, in *Planetary Nebulae in our Galaxy and Beyond*, IAU Symp.#234, Eds. Michael J. Barlow and Roberto H. Méndez, pp. 357
- Bakker, E.J., Waters, L.B.F.M., Lamers, H.J.G.L.M. et al. 1996, *A&A*, 310, 893
- Blöcker, T., Herwig, F., Driebe, T., 2000, *MmSAI*, 71, 711
- Bobrowsky, M., Sahu, K.C., Parthasarathy, M., García–Lario, P., 1998, *Nature*, 392, 469
- Conlon, E.S., Dufton, P.L., Keenan, F.P., & McCausland, R.J.H., 1991 *MNRAS*, 248, 82
- García–Lario, P., Manchado, A., Pych, W., Pottasch, S.R. 1997a, *A&AS*, 126, 479
- García–Lario, P., Parthasarathy, M., de Martino, D. et al. 1997b, *A&A*, 326, 1103
- Gauba, G. & Parthasarathy, M. 2004, *A&A*, 417, 201
- Gauba, G. & Parthasarathy, M. 2003, *A&A*, 407, 1007
- Herbig, G.H. 1993, *ApJ*, 407, 142
- Hog, E. Fabricius, C., Makarov, V.V. et al. 2000, *A&A*, 355, L27
- Hubeny, I., Stefl, S., & Harmanec, P. 1985, *Bull. Astron. Inst. Czechosl.*, 36, 214
- Jasniewicz, G., de Laverny, P., Parthasarathy, M. et al. 2004, *A&A*, 423, 353
- Kilian, J. 1992, *A&A*, 262, 171
- Kelly, D.M. & Hrivnak, B.J. 2005, *ApJ*, 629, 1040
- Klochkova, V.G., Yushkin, M.V., Miroshnichenko, A.S. et al. 2002, *A&A*, 392, 143
- Lang, K.R. 1980, *Astrophysical Formulae*, second edition, Springer–Verlag
- Lang, K.R. 1992, *Astrophysical Data : Planets and Stars*, Springer–Verlag
- Leitherer, C. 1988, *ApJ*, 326, 356
- McCausland, R.J.H., Conlon, E.S., Dufton, P.L., Keenan, F.P. 1992, *ApJ*, 394, 298
- Moehler, S. & Heber, U. 1998, *A&A*, 335, 985
- Moehler, S., Heber, U., Lemke, M., Napiwotzki, R. 1998, *A&A*, 339, 537
- Mooney, C.J., Rolleston, W.R.J., Keenan, F.P. et al. 2004, *A&A*, 419, 1123
- Moore, C.E. 1945, *A multiplet table of astrophysical interest*
- Omont, A., Loup, C., Forveille, T. et al. 1993, *A&A*, 267, 515
- Osterbrock, D.E., Fulbright, J.P., Martel, A.R. et al. 1996, *PASP*, 108, 277
- Parthasarathy, M., Pottasch, S.R. 1986, *A&A*, 154, L16
- Parthasarathy, M., Pottasch, S.R. 1989, *A&A*, 225, 521
- Parthasarathy, M., 1993, *A&A*, 414, L109
- Parthasarathy, M., García–Lario, P., Pottasch, S.R. et al. 1993 *A&A*, 267, L19
- Parthasarathy, M., García–Lario, P., de Martino, D. et al. 1995, *A&A*, 300, L25
- Parthasarathy, M., Vijapurkar, J., Drilling, J.S. 2000a, *A&AS*, 145, 269
- Parthasarathy, M., García–Lario, P., Sivarani, T. et al. 2000b, *A&A*, 357, 241
- Pottasch, S.R., Parthasarathy, M. 1988, *A&A*, 192, 182
- Ryans, R.S.I., Dufton, P.L., Mooney, C.J., et al. 2003, *A&A*, 401, 1119
- Sarkar, G., Parthasarathy, M., Reddy, B.E. 2005, *A&A*, 431, 1007
- Sahai, R., Sánchez Contreras, C., Morris, M. 2005, *ApJ*, 620, 948
- Schlegel, D.J., Finkbeiner, D.P., & Davis, M. 1998, *ApJ*, 500, 525
- Schmidt–Kaler, Th. 1982, in *Landolt–Börnstein: Numerical Data and Functional Relationships in Science and Technology*, vol. 2b, ed. K. Schaifers & H. H. Voigt (Berlin: Springer–Verlag)
- Thompson, H.M.A., Keenan, F.P., Dufton, P.L. et al. 2006, *MNRAS*, 368, 1749
- Volk, K., Hrivnak, B.J., Kwok, S. 2004, *ApJ*, 616, 1181
- Wackerling, L.R. 1970, *Mem. Roy. Astron. Soc.*, 73, 153
- Welch, C.A., Frank, A., Pipher, J.L., Forrest, W.J., 1999, *ApJ*, 522, L69
- Wood, P.R., Zarro, D.M. 1981, *ApJ*, 247, 247

Table 3. Emission lines in IRAS 22023+5249

$\lambda_{\text{obs.}}$ (Å)	$\lambda_{\text{lab.}}$ (Å)	Ident.	W_λ (Å)	$\log(gf)$	χ (eV)	$ \Delta\lambda $ (Å)	V_r km s ⁻¹
6043.111	6046.233	OI(22)	0.0968	-1.895	10.98–13.03		blend
	+ 6046.438			-1.675	10.98–13.03		
6092.139	6095.290	CII(24)	0.0131	-0.029	22.55–24.58	3.151	-140.22
6095.324	6098.510	CII(24)	0.0344	0.226	22.56–24.59	3.186	-141.86
6146.784	6150.10	FeII(46)	0.0292		3.21–5.21		blend
6148.099		UN	0.0828				
6296.769	6300.23	[OI](1F)	0.2832			3.461	-149.93
6337.276	6340.58	NII(46)	0.0479	-0.192	23.23–25.18	3.304	-141.46
6343.686	6346.86	NII(46)	0.5132	-0.901	23.22–25.18		blend
	+ 6347.109	SiII(2)		0.297	8.12–10.07		
6353.584	6357.0	NII(46)	0.0511		23.23–25.18	3.416	-146.34
6360.279	6363.88	[OI](1F)	0.0677			3.601	-154.89
6367.944	6371.371	SiII(2)	0.1989	-0.003	8.12–10.06	3.427	-146.49
6458.428		UN	0.1059				
6544.492	6548.1	[NII](1F)	2.817			3.608	-150.43
6579.838	6583.6 ^c	[NII](1F)	8.534				blend
6712.772	6717.0	[SII](2F)	0.5017			4.228	-173.96
6727.154	6731.3	[SII](2F)	1.007			4.146	-169.91
6848.05	6851.634	FeI(34)	0.1321	-5.320	1.61–3.41	3.584	-142.06
6998.28	7001.93	OI(21)	0.0786		10.94–12.70		blend
	+ 7002.22	OI(21)			10.94–12.70		
	7231.330 ^d	CII(3)		0.043	16.32–18.03		blend
	7236.420 ^d	CII(3)		0.299	16.32–18.03		blend
	7316.282 ^d	atmos.					
7373.821		UN	0.1893				
7458.676		UN	0.1119				
7462.426		UN	0.1229				
7464.227		UN	0.122				
7892.292		UN	0.134				
7993.283	7993.332	atmos.	0.0534				
8218.700	8223.128 ^d	NI(2)		-0.390	10.32–11.83		blend
8237.897	8242.389 ^d	NI(2)		-0.380	10.33–11.83		blend
8283.265		UN					blend
8441.864	8446.359	OI(4)	3.097	0.170	9.51–10.98		blend
	+ 8446.758	OI(4)		-0.050	9.51–10.98		
8745.796 ^d		UN					
8857.744 ^d		UN					

^c[NII](1F) 6583.6Å is blended with the emission component of CII(2) 6582.88Å P-Cygni profile. ^dThese emission lines are weak and are blended with the atmospheric absorption lines in this region.

Table 4. P–Cygni lines in IRAS22023+5249. Equivalent widths of the absorption and emission components of the P–Cygni profiles are given. Wind velocities are estimated from the blue absorption edges of the P–Cygni profiles.

λ_{lab} (Å)	Ident.	W_{λ} (absorption) (Å)	W_{λ} (emission) (Å)	log (gf)	χ (eV)	Wind Velocity km s ^{−1}
4340.462	H $_{\gamma}$	0.4908	1.915	−0.447	10.19–13.04	−187.81
4419.596 ^a	FeIII(4)	0.0991	0.0987	−2.218	8.23–11.04	blend
4431.019	FeIII(4)	0.0429	0.0506	−2.572	8.24–11.04	weak
4471.477	HeI(14)	0.3398	0.0828		20.95–23.72	blend
+4471.682	HeI(14)			−0.898	20.95–23.72	
4861.323 ^b	H $_{\beta}$	0.1235	6.779	−0.020	10.19–12.74	−192.63
5015.678	HeI(4)	0.3955	0.5047	−0.820	20.60–23.07	−185.39
5073.903	FeIII(5)	0.1361	0.0426	−2.557	8.65–11.09	weak
5086.701	FeIII(5)	0.0319	0.0338	−2.590	8.65–11.09	weak
5127.387	FeIII(5)	0.1048	0.1684	−2.218	8.65–11.07	−149.39
5156.111	FeIII(5)	0.1105	0.1964	−2.018	8.64–11.04	−149.14
5875.618	HeI(11)	0.5500	1.8790		20.87–22.97	blend
+ 5875.650	HeI(11)				20.87–22.97	
+ 5875.989	HeI(11)				20.87–22.97	
6562.797 ^c	H $_{\alpha}$	0.0443	37.86	0.710	10.19–12.08	−185.36
6578.050	CII(2)	0.1977	0.2779	−0.026	14.44–16.32	−191.55
6582.880 ^c	CII(2)			−0.327	14.43–16.32	blend
6678.154	HeI(46)	0.6693	0.8538	0.329	21.20–23.06	−182.16
7065.176	HeI(10)	0.3406	1.3600	−0.460	20.95–22.70	blend
+ 7065.707	HeI(10)			−1.160	20.95–22.70	

^aThe absorption component of FeIII(114) 4419.596Å is blended with OII(5) 4416.975 absorption feature. ^bThe emission components of the H $_{\beta}$ and H $_{\alpha}$ profiles have broad wings. Gaussian fits to the absorption and emission components of these profiles could not be obtained. Using IRAF, the equivalent widths of these components were estimated by subtracting the linear continuum between the points of interest and summing the pixels with partial pixels at the ends. ^c CII(2) 6582.88Å P–Cygni profile is blended with [NII](1F) 6583.6Å.

Table 5. Absorption (a) and emission (e) components of Na I D₂ (5889.953 Å) and Na I D₁ (5895.923 Å) lines in the spectrum of IRAS22023+5249 (LSIII +5224). W_{λ} are the equivalent widths of the components and V_r are the respective heliocentric radial velocities.

Component		IRAS22023+5249					
		Na I D ₂			Na I D ₁		
		$\lambda_{\text{obs.}}$ (Å)	W_{λ} (Å)	V_r (km s ^{−1})	$\lambda_{\text{obs.}}$ (Å)	W_{λ} (Å)	V_r (km s ^{−1})
1.	a	5886.341	0.0559	−169.10	5892.323	0.0529	−168.31
2.	e	5886.811	0.1314	−145.17	5892.763	0.0277	−145.92
3.	a	5888.530	0.1395	−57.61	5894.500	0.1355	−57.54
4.	a	5888.903	0.5054	−38.61	5894.881	0.3684	−38.15
5.	a	5889.500	0.6384	−8.20	5895.455	0.6241	−8.94
6.	e	5889.918	0.1221	+13.09	5895.902	0.1471	+13.8

Table 6a. Derived abundances for I22023. The values are for $\log \epsilon(\text{H}) = 12.0$. n refers to the number of lines of each species used for the abundance determination. For comparison we have listed the solar abundances ($\log \epsilon(\text{X})_{\odot}$) and average abundances for main sequence B–stars from the Ori OB1 association (Kilian, 1992).

X	IRAS22023+5249 ($T_{\text{eff}}=24000 \text{ K}$, $\log g=3.0$, $\xi_t=7 \text{ km s}^{-1}$)				$\log \epsilon(\text{X})_{\odot}$	Main sequence B–stars, Ori OB1 $\log \epsilon(\text{X})$
	n	$\log \epsilon(\text{X})$	σ	$[\text{X}/\text{H}]$		
He I	1	11.04 [†]	–	+0.11	10.93	11.04
C II	4	8.58 [†]	–	+0.19	8.39	8.23
N II	7	8.36	0.33	+0.44	7.92	7.72
O II	18	8.90	0.42	+0.21	8.69	8.60
Ne I	2	9.02	0.14	+0.94	8.08	–
Al III	1	< 6.79 [†]			6.47	–
Si III	3	7.43	0.59	–0.12	7.55	–

The abundances were derived using Kurucz’s WIDTH9 program.

[†]These values were derived from spectrum synthesis analysis using the SYNSPEC code.


Table 6b. Uncertainties in the derived abundances, $\Delta \log \epsilon(\text{X})$, due to uncertainties in the model atmospheric parameters

Element	ΔT_{eff} –1000K	$\Delta \log g$ +0.5	$\Delta \xi_t$ +1 km s ^{–1}	σ_m
N	–0.14	–0.18	–0.05	0.23
O	+0.05	+0.08	–0.08	0.12
Ne	–0.15	–0.26	–0.01	0.30
Si	–0.04	–0.06	–0.07	0.10

Table 7. Expansion velocities

Ident.	λ_{lab} Å	FWHM Å	V_{exp} km s ^{–1}
6300.23	[OI](1F)	1.334	31.76
6363.88	[OI](1F)	1.470	34.65
6548.1	[NII](1F)	0.844	19.34
6717.0	[SII](2F)	0.747	16.68
6731.3	[SII](2F)	0.739	16.47

[NII](1F) 6583.6Å line is blended with the emission component of CII(2) 6582.88Å P-Cygni profile and has not been used to estimate the expansion velocity.

High resolution optical spectrum of IRAS 22023+5249**Fig. .1.** Optical spectrum of IRAS22023+5249 (LS III +52°24)




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...




Fig. 1. Optical spectrum of IRAS22023+5249 (LS III +52°24) contd...